

# Package ‘metabolic’

October 10, 2023

**Title** Datasets and Functions for Reproducing Meta-Analyses

**Version** 0.1.2

**Description** Dataset and functions from the meta-analysis published in *Medicine & Science in Sports & Exercise*. It contains all the data and functions to reproduce the analysis. “Effectiveness of HIIE versus MICT in Improving Cardiometabolic Risk Factors in Health and Disease: A Meta-analysis”. Felipe Mattioni Maturana, Peter Martus, Stephan Zipfel, Andreas M Nieß (2020) <[doi:10.1249/MSS.0000000000002506](https://doi.org/10.1249/MSS.0000000000002506)>.

**License** CC0

**Encoding** UTF-8

**LazyData** true

**RoxygenNote** 7.2.1

**URL** <https://github.com/fmmattioni/metabolic>

**BugReports** <https://github.com/fmmattioni/metabolic/issues>

**Imports** tibble, magrittr, usethis, dplyr (>= 1.0.0), ggplot2 (>= 3.2.1), ggfittext, cli (>= 2.0.1), forcats, ggimage, patchwork (>= 1.0.0), scales, stringr, tidyr (>= 1.0.2), purrr, meta (>= 4.11-0), glue, rmarkdown

**Depends** R (>= 3.2)

**Suggests** knitr, here, Rd2roxygen, kableExtra, fansi, downloadthis (>= 0.2.0), spelling

**Language** en-US

**NeedsCompilation** no

**Author** Felipe Mattioni Maturana [aut, cre, cph]  
(<<https://orcid.org/0000-0002-4221-6104>>)

**Maintainer** Felipe Mattioni Maturana <[felipe.mattioni@med.uni-tuebingen.de](mailto:felipe.mattioni@med.uni-tuebingen.de)>

**Repository** CRAN

**Date/Publication** 2023-10-10 07:40:02 UTC

## R topics documented:

build_gofer . . . . .	2
build_report . . . . .	3
metabolic_gofer . . . . .	4
metabolic_meta . . . . .	6
perform_bind . . . . .	7
perform_meta . . . . .	8
plot_metabolic . . . . .	9
plot_small_study_effects . . . . .	10
read_paper . . . . .	11

<b>Index</b>	<b>12</b>
--------------	-----------

---

build_gofer	<i>Build a GOfER diagram (Graphical Overview for Evidence Reviews)</i>
-------------	------------------------------------------------------------------------

---

### Description

It is recommended to set `save = TRUE` and indicate the path to save to, as the plot is not going to look good in the Plots panel.

### Usage

```
build_gofer(page = c("1", "2"), save = FALSE, path, format = ".png")
```

### Arguments

page	A text string to indicate the page you would like to display. This GOfER has two pages (28 studies in page 1 and 28 studies in page 2).
save	A boolean to indicate whether to save the plot to disk.
path	Path to write to. It has to be a character string indicating the path and file name (without the extension). For example, <code>~/Documents/metabolic_gofer_page_1</code> will save <code>metabolic_gofer_page_1.png</code> to the Documents folder.
format	The file extension that you want to save the plot to. Only <code>.png</code> , is supported.

### Value

A patchwork object

### Examples

```
if (interactive()) {
  build_gofer(page = "1", save = TRUE, path = tempfile())
}
```

---

build_report	<i>Build HTML report</i>
--------------	--------------------------

---

## Description

Build an HTML report with all the results from the chosen clinical endpoint

## Usage

```
build_report(  
  endpoint = c("VO2max", "Flow-mediated Dilation", "BMI", "Body Mass", "Body Fat",  
    "Systolic Blood Pressure", "Diastolic Blood Pressure", "HDL", "LDL", "Triglycerides",  
    "Total Cholesterol", "C-reactive Protein", "Fasting Insulin", "Fasting Glucose",  
    "HbA1c", "HOMA-IR"),  
  path,  
  format = ".html"  
)
```

## Arguments

endpoint	The clinical endpoint to build the HTML report.
path	Path to write to. It has to be a character string indicating the path and file name (without the extension). For example, ~/Documents/metabolic_report will save metabolic_report.html to the Documents folder.
format	The file extension that you want to build the report with. Only .html, is supported.

## Value

an HTML file.

## Examples

```
if(interactive()) {  
  # Build an HTML report on VO2max  
  build_report(endpoint = "VO2max", path = tempfile())  
}
```

---

metabolic_gofer	<i>Dataset for building a GOfER diagram (Graphical Overview for Evidence Reviews)</i>
-----------------	---------------------------------------------------------------------------------------

---

## Description

A dataset containing the summary of the studies included in the meta-analysis. This dataset is used to build a GOfER with 'ggplot2' and 'patchwork'.

## Usage

```
metabolic_gofer
```

## Format

A data frame with 115 rows and 33 variables:

**study** last name of first author and year of publication

**groups** group allocated in the study, it may be either: HIIT (High-intensity Interval Training), SIT (Sprint Interval Training), or MICT (Moderate-intensity Continuous Training)

**sample\_population** population category from the study, it may be either: Healthy, Overweight/obese, Cardiac Rehabilitation, Metabolic Syndrome, or T2D (Type-2 Diabetes)

**sample\_fitness** the general fitness condition of the sample reported in the study, it may be either: Active, Sedentary, or N/R (Not Reported)

**sample\_men\_ratio** the men ratio (total men divided by sample size) reported in the study

**anamnese\_smoker** information whether participants in the sample were smokers, it may either: Y (Yes), N (No), or N/R (Not Reported)

**anamnese\_medicines\_to\_control\_BP** information whether participants in the sample were taking regular medication to control blood pressure, it may either: Y (Yes), N (No), or N/R (Not Reported)

**age** the age of each group reported in the study, in years

**design\_type\_of\_exercise** the type of exercise used for exercise training, it may be either running or cycling

**design\_sample\_size** the sample size of each group in the study

**design\_training\_duration** the training duration, in weeks

**design\_training\_frequency** the training frequency for each group used in the study

**design\_exercise\_intensity** the prescribed exercise intensity for each group

**hiie\_n\_reps** number of repetitions prescribed for the HIIE (High-intensity Interval Exercise) protocol

**hiie\_rep\_duration** length of repetitions prescribed for the HIIE (High-intensity Interval Exercise) protocol

**hiie\_work\_rest\_ratio** the work-rest ratio in the HIIE (High-intensity Interval Exercise) protocol

- compliance** compliance reported in each group and study
- endpoints\_vo2max** information on whether the reported p-value was significant comparing the effects pre- and post-training on VO<sub>2</sub>max (maximal oxygen uptake). If Yes, the reported p-value was less than 0.05; if No, the reported p-value was greater than 0.05
- endpoints\_fmd** information on whether the reported p-value was significant comparing the effects pre- and post-training on Flow-mediated Dilation. If Yes, the reported p-value was less than 0.05; if No, the reported p-value was greater than 0.05
- endpoints\_body\_mass** information on whether the reported p-value was significant comparing the effects pre- and post-training on Body Mass. If Yes, the reported p-value was less than 0.05; if No, the reported p-value was greater than 0.05
- endpoints\_body\_fat** information on whether the reported p-value was significant comparing the effects pre- and post-training on Body Fat. If Yes, the reported p-value was less than 0.05; if No, the reported p-value was greater than 0.05
- endpoints\_sbp** information on whether the reported p-value was significant comparing the effects pre- and post-training on Systolic Blood Pressure. If Yes, the reported p-value was less than 0.05; if No, the reported p-value was greater than 0.05
- endpoints\_dbp** information on whether the reported p-value was significant comparing the effects pre- and post-training on Diastolic Blood Pressure. If Yes, the reported p-value was less than 0.05; if No, the reported p-value was greater than 0.05
- endpoints\_hdl** information on whether the reported p-value was significant comparing the effects pre- and post-training on HDL. If Yes, the reported p-value was less than 0.05; if No, the reported p-value was greater than 0.05
- endpoints\_ldl** information on whether the reported p-value was significant comparing the effects pre- and post-training on LDL. If Yes, the reported p-value was less than 0.05; if No, the reported p-value was greater than 0.05
- endpoints\_triglycerides** information on whether the reported p-value was significant comparing the effects pre- and post-training on Triglycerides. If Yes, the reported p-value was less than 0.05; if No, the reported p-value was greater than 0.05
- endpoints\_total\_cholesterol** information on whether the reported p-value was significant comparing the effects pre- and post-training on Total Cholesterol. If Yes, the reported p-value was less than 0.05; if No, the reported p-value was greater than 0.05
- endpoints\_insulin** information on whether the reported p-value was significant comparing the effects pre- and post-training on Fasting Insulin. If Yes, the reported p-value was less than 0.05; if No, the reported p-value was greater than 0.05
- endpoints\_glucose** information on whether the reported p-value was significant comparing the effects pre- and post-training on Fasting Glucose. If Yes, the reported p-value was less than 0.05; if No, the reported p-value was greater than 0.05
- endpoints\_homa** information on whether the reported p-value was significant comparing the effects pre- and post-training on HOMA-IR (insulin resistance). If Yes, the reported p-value was less than 0.05; if No, the reported p-value was greater than 0.05
- endpoints\_bmi** information on whether the reported p-value was significant comparing the effects pre- and post-training on BMI (body mass index). If Yes, the reported p-value was less than 0.05; if No, the reported p-value was greater than 0.05

**endpoints\_crp** information on whether the reported p-value was significant comparing the effects pre- and post-training on C-reactive Protein. If Yes, the reported p-value was less than 0.05; if No, the reported p-value was greater than 0.05

**endpoints\_hba1c** information on whether the reported p-value was significant comparing the effects pre- and post-training on HbA1c (glycated hemoglobin). If Yes, the reported p-value was less than 0.05; if No, the reported p-value was greater than 0.05

### Source

[https://journals.lww.com/acsm-msse/Fulltext/2021/03000/Effectiveness\\_of\\_HIIE\\_versus\\_MICT\\_in\\_Improving.12.aspx](https://journals.lww.com/acsm-msse/Fulltext/2021/03000/Effectiveness_of_HIIE_versus_MICT_in_Improving.12.aspx)

---

metabolic_meta	<i>Dataset for reproducing the meta-analysis</i>
----------------	--------------------------------------------------

---

### Description

A dataset containing the processed data from the studies necessary to reproduce the meta-analysis.

### Usage

metabolic\_meta

### Format

A data frame with 391 rows and 21 variables:

**study** last name of first author and year of publication

**endpoint** the clinical endpoint analyzed, it may be either: VO2max (maximal oxygen uptake), Flow-mediated Dilation, BMI (body mass index), Body Mass, Body Fat, Systolic Blood Pressure, Diastolic Blood Pressure, HDL, LDL, Triglycerides, Total Cholesterol, C-reactive Protein, Fasting Insulin, Fasting Glucose, HbA1c (glycated hemoglobin), or HOMA-IR (insulin resistance)

**population** population category from the study, it may be either: Healthy, Overweight/obese, Cardiac Rehabilitation, Metabolic Syndrome, or T2D

**age** the median age between the groups in the study, in years

**category\_age** age category based on the age column, it may be either: < 30 y, 30 - 50 y, or > 50 y

**duration** the training duration, in weeks

**category\_duration** training duration category based on the duration column, it may be either: < 5 weeks, 5 - 10 weeks, or > 10 weeks

**men\_ratio** the men ratio (total men divided by sample size) reported in the study

**category\_men\_ratio** men ratio category based on the men\_ratio column, it may be either: < 0.5 or > 0.5

**type\_exercise** the type of exercise used for exercise training, it may be either running or cycling

- bsln** the baseline value reported for the clinical endpoint (the median between groups is used)
- bsln\_adjusted** the adjusted baseline value for the clinical endpoint. Values were adjusted according to their categories described in the paper. For example, VO2max values were adjusted to their age and sex percentile ranks, etc. From these values, the categories are defined in 'category\_bsln'
- category\_bsln** the baseline category based on the bsln column
- N\_HIIE** sample size of the HIIE (High-intensity Interval Exercise) group
- Mean\_HIIE** mean difference between pre- and post-training in the HIIE (High-intensity Interval Exercise) group
- SD\_HIIE** standard deviation of the difference between pre- and post-training in the HIIE (High-intensity Interval Exercise) group
- N\_MICT** sample size of the MICT (Moderate-intensity Continuous Training) group
- Mean\_MICT** mean difference between pre- and post-training in the MICT (Moderate-intensity Continuous Training) group
- SD\_MICT** standard deviation of the difference between pre- and post-training in the MICT (Moderate-intensity Continuous Training) group
- HIIE** the type of HIIE used in the study: it may be either: HIIT (High-intensity Interval Training) or SIT (Sprint Interval Training)
- desired\_effect** the desired effect expected for post-training improvements. This is needed simply to display the effects related to HIIE and MICT on the same side of the forest plot throughout the clinical endpoints

### Source

[https://journals.lww.com/acsm-msse/Fulltext/2021/03000/Effectiveness\\_of\\_HIIE\\_versus\\_MICT\\_in\\_Improving.12.aspx](https://journals.lww.com/acsm-msse/Fulltext/2021/03000/Effectiveness_of_HIIE_versus_MICT_in_Improving.12.aspx)

---

perform\_bind

*Combine the subgroup meta-analyses*

---

### Description

Combine the subgroup meta-analyses to ...

### Usage

```
perform_bind(x)
```

### Arguments

x An object retrieved from [perform\\_meta](#).

### Value

a [tibble](#) with named lists.

**Examples**

```

if (interactive()) {
# Perform meta-analysis on VO2max
results <- perform_meta(endpoint = "VO2max")
results

# Combine Overall and Subgroups meta-analysis results
results_bind <- perform_bind(results$meta_analysis)
results_bind
}

```

---

perform\_meta

*Perform meta-analysis*


---

**Description**

Perform the meta-analysis, sensitivity analysis, and meta-regression on the chosen clinical endpoint.

**Usage**

```

perform_meta(
  endpoint = c("VO2max", "Flow-mediated Dilation", "BMI", "Body Mass", "Body Fat",
    "Systolic Blood Pressure", "Diastolic Blood Pressure", "HDL", "LDL", "Triglycerides",
    "Total Cholesterol", "C-reactive Protein", "Fasting Insulin", "Fasting Glucose",
    "HbA1c", "HOMA-IR")
)

```

**Arguments**

endpoint      The clinical endpoint to perform the meta-analysis and meta-regression.

**Value**

a [tibble](#) with named lists.

**Examples**

```

if (interactive()) {
# Perform meta-analysis on VO2max
results <- perform_meta(endpoint = "VO2max")
results

# Access results of Overall meta-analysis
results$meta_analysis$Overall

# Access results of Age meta-regression
results$meta_regression$Age
}

```



---

plot_metabolic	<i>Plot results</i>
----------------	---------------------

---

### Description

Plot results from the `perform_meta()` and `perform_bind()` function. Please, see 'Details' and 'Examples'.

### Usage

```
plot_metabolic(x, save = FALSE, path, format = ".png")
```

### Arguments

<code>x</code>	an object obtained from the meta-analysis results. See 'Details'.
<code>save</code>	A boolean to indicate whether to save the plot to disk.
<code>path</code>	Path to write to. It has to be a character string indicating the path and file name (without the extension). For example, <code>~/Documents/metabolic_gofer_page_1</code> will save <code>metabolic_gofer_page_1.png</code> to the Documents folder.
<code>format</code>	The file extension that you want to save the plot to. Only <code>.png</code> , is supported.

### Details

This function can be used to plot the results derived from both `perform_meta()` and `perform_bind()`. It can produce forests and bubble plots, depending on the object passed to the function.

### Value

a plot.

### Examples

```
if(interactive()) {  
  # Perform meta-analysis on V02max  
  results <- perform_meta(endpoint = "V02max")  
  
  # Plot Overall meta-analysis results  
  results$meta_analysis$Overall %>%  
    plot_metabolic()  
  
  # Plot Age meta-regression results  
  results$meta_regression$Age %>%  
    plot_metabolic()  
  
  # Plot overview of Overall and Subgroups meta-analysis results  
  results_bind <- perform_bind(results$meta_analysis)  
  results_bind %>%  
    plot_metabolic()  
}
```

```
# Plot sensitivity analysis results
results$sensitivity_analysis$Overall %>%
  plot_metabolic()
}
```

---

plot\_small\_study\_effects

*Plot small-study effects analysis*

---

## Description

Plot small-study effects analysis

## Usage

```
plot_small_study_effects(x, save = FALSE, path, format = ".png")
```

## Arguments

x	an object of class meta
save	A boolean to indicate whether to save the plot to disk.
path	Path to write to. It has to be a character string indicating the path and file name (without the extension). For example, ~/Documents/small_study_effects will save small_study_effects.png to the Documents folder.
format	The file extension that you want to save the plot to. Only .png, is supported.

## Value

a plot.

## Examples

```
## Not run:
# Perform meta-analysis on V02max
results <- perform_meta(endpoint = "V02max")

# Plot small-study effects results
results$meta_analysis$Overall %>%
  plot_small_study_effects()

## End(Not run)
```

---

<code>read_paper</code>	<i>Read the paper</i>
-------------------------	-----------------------

---

**Description**

This function will open the published paper in the journal website for you to read it in your default browser.

**Usage**

```
read_paper()
```

**Examples**

```
read_paper()
```

# Index

## \* datasets

metabolic\_gofer, 4

metabolic\_meta, 6

build\_gofer, 2

build\_report, 3

metabolic\_gofer, 4

metabolic\_meta, 6

perform\_bind, 7

perform\_meta, 7, 8

plot\_metabolic, 9

plot\_small\_study\_effects, 10

read\_paper, 11

tibble, 7, 8